

spira breviter conoidea, apice obtusula; anfr. 5 convexi, ultimus inflatus, subangulatus, antice descendens, subtus nitens, viridulo-flavus, circa umbilicum mediocrem, pervium subcompressus; apertura diagonalis, ampliter lunato-rotundata, intus livide margaritacea; peristoma albidum, marginibus callo tenui, nitido junctis, supero leviter dilatato, non reflexo, columellari parum obliquo, superne dilatato, fornicatim reflexo, umbilici partem occultante. — Diam. maj. 24, min. 21, alt. 16 mill.

VAR. β minor, peristomate violacescente, umbilico vix partim oblecto, brunneo circumdato. — Diam. maj. 20, min. 17, alt. 13 mill.

Hab. in Oceania (coll. Crosse).

Species præcedenti peraffinis, sed minor, testa tenui, validius rugoso-striata, nec subgranosa, subtus nitente, columella obliqua, apertura subampla et umbilico facile distinguitur.

H. C.

Description de **coquilles inédites**,

PAR A. MORELET.

I. AFRIQUE.

1. BULIMUS INGENUUS.

T. vix perforata, ovato-conica, tenuis, subdiaphana, subtiliter striatula et sub-lente minutissime decussata, nitida, albido-straminea, fasciis badio-rufis in anfr. prioribus interruptis, in ultimo continuis, ornata; spira co-

nica, apice acuta, nitide rufa; anfr. 6 convexiusculi, ultimus ventricosus, spira paulo minor; columella recta; apertura semi-ovalis, intus vivide fasciata, marginibus tenuibus, rectis, columellari superne in lamellam triangularem parvulam dilatato, revoluto. — Longit. 17, diam. 10 mill.

Hab. in insula Mayotte.

2. BULIMUS SPINULA.

T. subperforata, turrata, tenuis, arcuatim striolata, diaphana, virenti-cornea; spira elongata, apice obtusa; anfr. 6-7 subcylindraceuti, mediocriter convexi, sutura profunda minute denticulata conjuncti, ultimus spiræ dimidium æquans; apertura truncato-ovalis, marginibus tenuibus, rectis, columellari leviter dilatato, reflexiusculo. — Longit. 7-8, diam. 2-2 2/5 mill.

Hab. ad ripas fluv. Gabon.

3. MELANIA PSORICA.

T. cylindraceuto-turrata, truncata, solida, striata, pallide olivacea; anfr. superst. 4 plano-convexi, sutura profunda constricti, costis longitudinalibus, in medio anfractus ultimi evanescentibus, et liris paucis decurrentibus, tuberculoso-rugata, basi spiraliter costulata; apertura ovato-acuminata, parva. — Longit. 18, diam. 6 mill.

Hab. Madagascar.

4. MELANIA SORICULATA.

T. pyramidalis-turrata, crassiuscula, lutea vel flava, infra suturas et basi nigro vel rubido late fasciata; spira sensim attenuata, apice truncata; anfr. 9-10 planati, spiraliter costulati, priores plicis longitudinalibus costisque decurrentibus eximie clathrati; apertura parva, ovato-

acuta, basi subeffusa, intus fasciata, margine columellari declivi, calloso. — Longit. 18, diam. 6 mill.

Hab. ad Grand Bassam Africa æquinoxialis.

5. NERITINA CRISTATA.

T. semi-globosa, patula, apice breviter conica, sæpius erosa, capillaceo-striata, sub indumento nigro castanea maculisque luteis, squammæformibus, aliquando seriatis, in specim. adultis parum conspicuis, ornata; anfr. 3 convexiusculi, celeriter crescentes, ultimus permagnus, involvens, plicam prominentem, cristatam, superne canaliculatam, suturam ab anfractus dimidio usque ad basim concomitantem, exhibens; apertura late semi-ovalis, auriculata, canaliculata, intus cærulescens; callum columellare convexum, declive, carneo-violaceum, margine medio minute denticulato. — Operculum crassum, radiatum, carneum, ad nucleum violaceo diffuse maculatum. — Longit. 21, diam. 20 mill.

Hab. in flumine Como Gabonensi.

II. INDO-CHINE.

6. PALUDOMUS CYANOSTOMUS.

T. parva, oblonga, crassa, apice attenuata, sub indumento nigro tenacissimo albido-cornea, nitida, exitissime decussatula; anfr. 3 convexiusculi, ultimus ovatus, magnus, spiram fere æquans; apertura integra, ovalis, superne acuta, plumbeo-cyanea, marginibus callosis, columellari strictim dilatato, expresso. — Longit. 8, diam. 4 mill.

Hab. Siam.

7. UNIO MASSINI.

T. subrhomboidea, inæquilateralis, tumida, leviter

striata, antice attenuata, subrostrata, postice ovata; area in alam compressam, parvam, dilatata; margo superior rectilinearis, ascendens; basalis arcuatus; epidermis brunneo-virens, postice obscure radiata; umbones inflati, decorticati; margarita albo-cærulea, iridescens; laminæ cardinales compressæ, acutæ, arcuatæ; dentes lamelli-formes, marginales, minutim serrulatæ. — Altit. 42, latit. 58, diam. 51 mill.

Hab. in Cochinchina.

Differt ab U. abnormi statura præcipue et cardine arcuato, nec recto; etenim U. abnormis adultus usque ad 75 mill. altitudinis, 110 longitudinis et 67 diametri crescit.

III. AUSTRALIE.

8. HELIX SEMINIGRA.

T. subobtecte umbilicata, turbinata, solida, inæqualiter striata, nitida; spira conoideo-rotundata, apice obtusa; anfr. 6 1/2 vix convexi, priores fulvacei, obscure fasciati, ultimus atro-purpureus, celeriter descendens, basi subplanatus, circa umbilicum paululum excavatus; sutura irregulariter crenulata, pallide marginata; apertura perobliqua, ovato-lunaris, intus saturate plumbea; peristoma subincrassatum, album, marginibus callo tenui junctis, breviter expansis, columellari superne dilatato, patente, umbilicum parvum semitegente. — Diam. maj. 55, min. 28, altit. 21 mill.

Hab. Queen's Land Australiæ.

9. PUPINA COXI.

T. late rimata, pupæformis, cornea, subdiaphana, nitida, sublente confertim striatula; spira oblongo-

conica, apice obtusiuscula; anfr. 6 convexiusculi, ultimus vix striatus, antice subplanatus, angulis spiratibus obsoletissime notatus, circa rimam umbilicalem funiculatus, breviter descendens; apertura circularis, bicanaliculata; peristoma subincrassatum, expansum, pallidum, canalibus recte dissectum, margine externo sulco superficiali circulatim bipartito, columellari strictiore. — Operculum tenue, corneum, arctispirum, nucleo concavo. — Longit. 14, diam. 5 1/2 mill.

Hab. Port-Curtis Australiæ.

10. MELAMPUS TETRICUS.

T. obtecte subrimata, ovata, solida, sublente tenerime striata, epidermide atro-viridi, sæpe deficiente, vestita; spira brevis, conica, apice erosa; anfr. 5 subplani, sutura irregulari conjuncti, ultimus basi attenuatus, 3/5 longitudinis æquans; apertura anguste semi-ovalis, biplicata, intus atro-purpurea; plica parietalis 1 alba, compressa, subhorizontalis; columellaris torta, marginem attingens; perist. acutum, margine dextro medio callo leviter prominulo munito, columellari incrassato, reflexo, adnato. — Longit. 10, latit. 5 1/5 mill.

Hab. « Nouvelle-Galles du Sud. »

BIBLIOGRAPHIE.

Cours de **Paléontologie stratigraphique**, professé au muséum d'histoire naturelle par **A.**